

## BASELINE DOSSIER

### ***Bacillus subtilis QST 713*** Microbial pest control agent against plant pathogenic fungi and bacteria

Dossier according to OECD guidance for industry data submissions for microbial pest control products and their microbial pest control agents August 2006

Summary documentation, Tier II

Annex IIM, Section 4

#### **Point IIM 6: Metabolism and residue studies**

Date: July 2015

Applicant

Bayer CropScience AG

This document and/or any copy rights to rights of the owner of this document may fall under regulation and use of the owner of this document or its rights of this document may be prohibited and violate the rights of this document and its owner.

Furthermore, this document and/or any copy rights to rights of the owner of this document may fall under regulation and use of the owner of this document or its rights of this document may be prohibited and violate the rights of this document and its owner.

It may be subject to rights of the owner of this document and/or any copy rights to rights of the owner of this document may fall under regulation and use of the owner of this document or its rights of this document may be prohibited and violate the rights of this document and its owner.

Consequently, any commercial publication may fall under regulation and use of the owner of this document or its rights of this document may be prohibited and violate the rights of this document and its owner.

Without the permission of the owner of this document or its rights of this document may be prohibited and violate the rights of this document and its owner.

Applicant  
Bayer CropScience AG



M-535397-01-3

**Table of Contents**

IIM 6	Metabolism and residues studies on the microbial pest control agent.....	5
IIM 6.1	Rationale for waiver of residue data based on information showing that MPCA is not hazardous to mammals, i.e. lack of potential for a known mammalian toxin and negative result from the acute oral toxicity test.....	5
IIM 6.2	Rationale for waiver based on a substantiated estimation that MPCA is unlikely to occur on treated food/feed stuffs in concentrations considerably higher than under natural conditions.....	5
IIM 6.3	Persistence and likelihood of multiplication in or on crops, feedingstuffs or foodstuffs .....	5
IIM 6.4	Further information required .....	8
IIM 6.4.1	Non-viable residues.....	8
IIM 6.4.2	Viable residues.....	9
IIM 6.5	Summary of residue behaviour and overall evaluation.....	10

It may be subject and/or any is the property of Bayer AG  
Consequently, this copy rights to rights of its affiliates and intellectual property parties.  
any commercial publication may fall under a owner and third party data protection and/or publishing and  
without the permission distribution and use of this document or its contents therefore  
exploitation, distribution and use, reproduction data protection and/or publishing and  
be prohibited and violate the rights of this document or its owner.

**OWNERSHIP STATEMENT**

This document, the data contained in it and copyright therein are owned by Bayer CropScience AG. No part of the document or any information contained therein may be disclosed to any third party without the prior written authorisation of Bayer CropScience AG.

It may be subject and/or any is the property of its affiliates such as intellectual owner and third parties. Furthermore, copy rights to rights of the property of Bayer AG without the permission of the owner of this document and use of this document and/or protection regime or its contents therefore. Consequently, any commercial publication may fall under a distribution and use of this document and protection regime and/or publishing regime and be prohibited and violate the rights of its owner. Consequently, any commercial publication may fall under a distribution and use of this document and protection regime and/or publishing regime and be prohibited and violate the rights of its owner.

**Introduction**

This document summarizes all data submitted for the initial evaluation of *Bacillus subtilis* QST 713 as an active substance under Directive 91/414. Data provided in the initial dossier and in subsequent additional submissions are listed chronologically under the respective data point according to the OECD dossier guidance (2006).

This document is further named as "**Baseline Dossier**" since it presents all data previously submitted.

It may be subject and/or any is the property of its affiliates such as affiliates of Bayer AG owner and intellectuates. without the permission, distribution under a regulatory parties. and protection and/or publication regime and contents therefore. Consequently, any commercial publication may fall under a third party rights of this document or its owner. and use of this document or its owner may therefore. be prohibited and violate the rights of its owner. and use of this document or its owner may therefore.

**IIM 6 Metabolism and residues studies on the microbial pest control agent****IIM 6.1 Rationale for waiver of residue data based on information showing that MPCAs are not hazardous to mammals, i.e. lack of potential for a known mammalian toxin and negative result from the acute oral toxicity test**

Studies were submitted in Annex II, Section 3, Points IIM 5.3.2 – 5.3.3 on acute oral and inhalatory toxicity in rats. The results showed the absence of any clinical signs and therefore it is concluded that *B. subtilis* poses no risk to mammals, and can be classified as non-toxic. Since *B. subtilis* QST713 is not infectious to mammals and does not produce metabolites toxic to other organisms, deposits of this microorganism after application of the plant protection product will not impose any health risk for consumers. Consequently no residues data are required.

**IIM 6.2 Rationale for waiver based on a substantiated estimation that MPCAs are unlikely to occur on treated food/feed stuffs in concentrations considerably higher than under natural conditions****Report:**

IIM 6.2/01; [REDACTED]; 1993; M-484952-01-1

**Title:**Systematics and ecology of *Bacillus***Report No.:**

M-484952-01-1

**Document No.:**

M-484952-01-1

**Guideline(s):**

not specified

**Guideline deviation(s):**

not specified

**GLP/GEP:**

no

Based on reported characteristics of *B. subtilis* species, as uniformly found in the scientific literature, under consideration of the envisaged application and relevant properties of strain QST713, we can conclude that populations on treated food/feed stuffs may be higher only directly after application, and will deplete significantly until the harvest of the treated crop. This refers also to Point IIM 6.3, which presents the data on the possibility of multiplication of the strain of interest on food stuffs and feeding stuffs.

It has to be considered that *B. subtilis* is a non-pathogenic, ubiquitous micro-organism, prevalent in the microflora of different environmental compartments and media, including the leaf surface and foodstuff ([REDACTED], 1997).

**IIM 6.3 Persistence and likelihood of multiplication in or on crops, feedingstuffs or foodstuffs****EU-Dossier: Doc M-11B, Point 6.1****Report:**

IIM 6.3/01; [REDACTED]; 1989; M-488048-01-1

**Title:**

Biocontrol on leaf surface

**Report No.:**

M-488048-01-1

**Document No.:**

M-488048-01-1

**Guideline(s):**

not specified

**Guideline deviation(s):**

not specified

**GLP/GEP:**

no

**Report:**

IIM 6.3/02; [REDACTED]; [REDACTED] C. D.; [REDACTED] R. L.; [REDACTED] B.

A.; [REDACTED] B.; [REDACTED] E. C.; [REDACTED] S.; [REDACTED] S. A.; [REDACTED] S. A.;

**Title:**

[REDACTED]; [REDACTED] E. O.; 1997; M-528163-01-1

Final decision document: TSCA section 5 (H) (4) exemption for *Bacillus subtilis***Report No.:**

M-528163-01-1

**Document No.:**

M-528163-01-1

**Guideline(s):**

--

**Guideline deviation(s):**

--

**GLP/GEP:**

no

**Report:**

KIIM 6.3/03; [REDACTED]; 2000; M-497595-01-1

Title:

Longevity study of Serenade (QST713) on pepper leaf surface in greenhouse conditions

Report No.:

M-497595-01-1

Document No.:

M-497595-01-1

Guideline(s):

not specified

Guideline deviation(s):

not specified

**GLP/GEP:**

no

**Report:**

KIIM 6.3/04; [REDACTED]; 1993; M-484952-01-1

Title:

Systematics and ecology of *Bacillus*

Report No.:

M-484952-01-1

Document No.:

M-484952-01-1

Guideline(s):

not specified

Guideline deviation(s):

not specified

**GLP/GEP:**

no

**Report:**

KIIM 6.3/05; [REDACTED]; [REDACTED]; [REDACTED]; 1993; M-528182-01-1

Title:

Biocontrol of postharvest diseases of apple using *Bacillus* spp. isolated from stored apples

Report No.:

M-528182-01-1

Document No.:

M-528182-01-1

Guideline(s):

--


Guideline deviation(s):

--

**GLP/GEP:**

no

Strain QST 713 of *B. subtilis* is intended to be applied onto the foliage. Regarding the intended fields of use residues of *B. subtilis* on leaf surfaces are associated with the establishment of colonization, the prime phenomenon of this contact-biofungicide and bactericide. Colonization of treated foliage provides a protective layer and basically is involved in the mode of action of *B. subtilis* against pathogen attack (see Fig. 6.3-01 and see Section 1, Point 2.3.2).



**Fig. 6.3-01: Grape powdery mildew spore on a dense “carpet” of *B. subtilis* cells on squash**

[REDACTED] (1989) addresses to the problems involved in foliar application of biocontrol agents, including adhesion of preparations to the mainly hydrophobic leaf surface and the usually unfavourable environmental conditions restricting microbial growth (and explaining the generally

slight occurrence of growing saprophytic bacteria on the leaf surface and the low covering percentage (<1%) of micro-organisms on the leaf surface of temperate plants). In addition, *B. subtilis* cells will stop growth after depletion of organic matter supply (EPA 1997), e.g. the fungal pathogen.

Therefore, the protective layer of *B. subtilis* cells colonizing the leaf surface will not maintain stability or persistence for long in this stressed micro-habitat.

The applicant conducted a study on longevity of QST 713 strain of *B. subtilis* on pepper leaf surface under greenhouse conditions, applied at a rate of 6.7 kg/ha formulated product (████████&████ 2000): an initial increase in colony forming units (CFU) was followed by a sharp decline in CFU-counts by day 5 to low levels of surviving cells persisting for ~ 3 weeks (Fig. 6.1-02).




Fig. 6.1-02: Time course of CFU-counts of *B. subtilis* strain QST 713 recovered from pepper leaf surface

To assess the survival of this strain in the field and in summertime, especially with an assumed negative effect of increased UV radiation, further studies under ambient field conditions are planned.

However, the unfavourable environmental conditions prevailing on the leaf surface will not impede the efficacy of QST 713 strain of *B. subtilis* since the preparation will be added several times at appropriate intervals and in sufficient amounts so that long-term survival or long-term colonization of the foliage is not necessitated – in fact for the induction of systemic resistance in the plant, presenting one mode of action (please refer to Annex IIM, Section 1, Point 2.3.2), an initial challenge by a non-pathogenic species may elicit the resistance response (████, 1989).

*B. subtilis* may form endospores under nutrient shortage and environmental stress. These endospores might possibly be dissipated via wind and water to other environmental compartments (████, 1993), where they may contribute to the existent population of *B. subtilis*, e.g.:

████ et al (1995) recovered 95 bacterial isolates from stored apples for screening of biocontrol activity and identified 7 among 30 *Bacillus* isolates as *B. subtilis* which acted effectively against molds.

With regard to *B. subtilis* cells remaining on harvested fruit the chance of proliferation during processing of raw products (grapes, apples to wine, juice respectively) is not relevant since a) in wine fermentation conditions are unfavourable and b) in juice production microbial contaminants are heat killed at a processing temperature of ~ 90°C), while conditions do not favour endospore formation.

In this context, it has to be considered that *B. subtilis* is a non-pathogenic, ubiquitous micro-organism, prevalent in the micro-flora of different environmental compartments and media, including the leaf surface and foodstuff (████, 1993).

**IIM 6.4 Further information required****IIM 6.4.1 Non-viable residues****Report:**

KIIM 6.4.1/01;

Title:

Final decision document: TSCA section 5 (H) (4) exemption for *Bacillus subtilis*

Report No.:

M-528163-01-1

Document No.:

M-528163-01-1

Guideline(s):

--

Guideline deviation(s):

--

GLP/GEP:

no

**Report:**

KIIM 6.4.1/02; [REDACTED], D : 2000; M-528847-01-1

Title:

Absence of pharmaceuticals in 713.

Report No.:

M-528847-01-1

Document No.:

M-528847-01-1

Guideline(s):

not specified

Guideline deviation(s):

not specified

GLP/GEP:

no

This refers to Annex II, Section 1, Point IIM 2.6. *B. subtilis* does not produce significant quantities of extracellular enzymes or toxins and is generally considered to have a low degree of virulence (EPA 1997). The results of the submitted toxicological studies on rodents demonstrate that QST 713 strain of *B. subtilis* does evidently not produce toxins (please refer to Annex II, Section 3, Point IIM 5.3.2 – 5.3.3).

In conclusion, non-viable residues originating from *B. subtilis* QST 713 are not considered to pose a risk to human health and the environment.

Included under 1<sup>st</sup> Additional Submission Amendment not included in monograph

Secondary metabolites of strain QST 713 of *B. subtilis* have been analyzed and determined to be in the known class of lipopeptides ([REDACTED], 2001, submitted in Doc JII). The complete chemical analysis of the strain QST 713 proved that no compounds are produced that are not well known in the literature. Based on a literature search it can be stated that the strain QST 713 does not produce metabolites that are used in human medicine ([REDACTED], 2000). Please refer to Annex II, Section 1, Point IIM 2.6.

Included under 2<sup>nd</sup> Additional Submission

*Bacillus subtilis* produces several different secondary metabolites. Detailed information on secondary metabolites formed by the strain QST 713 of *B. subtilis* has been submitted to all Member States in October 2001 for the evaluation for the Annex I inclusion ([REDACTED], 2001). Please refer to Annex II, Section 1, Point IIM 2.6.

Included under 3<sup>rd</sup> Additional Submission

Additional information was submitted in response to the EU Evaluation to provide further information on toxins produced by *B. subtilis* QST 713 (date of submission: June 2002). Please refer to Annex II, Section 1, Point IIM 2.6.

**IIM 6.4.2 Viable residues**

**Report:** KIIM 6.4.2/01; [REDACTED]; [REDACTED]; 1991; M-486912-01-1

Title: On the safety of *Bacillus subtilis* and *B. amyloliquefaciens*: a review

Report No.: M-486912-01-1

Document No.: M-486912-01-1

Guideline(s): not applicable

Guideline deviation(s): not applicable

GLP/GEP: no

**Report:** KIIM 6.4.2/02; [REDACTED]

Title: Final decision document: TSCA section 5 (H) (4) exemption for *Bacillus subtilis*

Report No.: M-528163-01-1

Document No.: M-528163-01-1

Guideline(s): --

Guideline deviation(s): --

GLP/GEP: no

**Report:** KIIM 6.4.2/03; [REDACTED]; 1992; M-484952-01,

Title: Systematics and ecology of *Bacillus*

Report No.: M-484952-01-1

Document No.: M-484952-01-1

Guideline(s): not specified

Guideline deviation(s): not specified

GLP/GEP: no

**Report:** KIIM 6.4.2/04; [REDACTED], J. 2001; M-487847-01-1

Title: Serenade WP residue on wine grapes

Report No.: 010147

Document No.: M-487847-01-1

Guideline(s): --

Guideline deviation(s): --

GLP/GEP: no

**EU Dossier Doc M-IIB, Point 6.2**

Due to the fact that the active ingredient is a viable micro-organism of ubiquitous occurrence and predominance in the soil-microflora the term residue is not applicable to this preparation.

Specifically no residue metabolism can be stated, since a micro-organism does not follow first order kinetics.

Residue studies were regarded as being dispensable for the above given and following additional reasons:

- With regard to its natural global distribution and non-pathogenic character *B. subtilis* cells left on the surface of treated areas or plant products do not imply health or environmental impacts (Bar & [REDACTED] 1991; EPA 1997).
- *B. subtilis* has been used for enzyme production on a large industrial scale, and is even used for food production without having caused health or environmental hazards or damages (e.g. [REDACTED], 1993).
- *B. subtilis* has no special attachment ability to plants or plant products, i.e. there is no compatibility comparable to host-pathogen interactions (EPA 1997).

- Colonization of the leaf surface by *B. subtilis* contributes largely to the protective effect against bacterial and fungal pathogens and according to the recommended frequent applications this protective layer may be renewed almost weekly.
- A plant product (fruit) carrying a layer built up of *B. subtilis* can easily be washed with water prior to consumption or juice production.

Included under 3<sup>rd</sup> Additional Submission

A relevant study which demonstrates the decline of *Bacillus subtilis* strain QST 713 on grapes after application of Serenade WP (███████, 2001) was submitted in June 2002 and is cited in the Addendum 1 to the Monograph (date of issue: 04.12.2002). According to this study *Bacillus subtilis* strain QST 713 declines on grapes after application of 10 kg Serenade WP/ha within four weeks after the last application to values below  $7.1 \times 10^3$  CFU/g berries.

**On the ECCO Working Group Evaluation Meeting on 26.03.2003, it was stated that no data are required for the time being as MRIs are not considered necessary.**

**IIM 6.5 Summary of residue behaviour and overall evaluation**

**EU-Dossier: Doc M-II B, Point 6.3**

Primarily the low health and environmental risk potential of *B. subtilis* and its ubiquitous distribution indicate that residual *B. subtilis* cells may present only a low risk potential.

Secondly, the unfavourable environmental conditions prevailing on the leaf surface and the dependence of *B. subtilis* on organic matter supply are restricting its growth, as shown in the submitted study report (please refer to Point IIM 6.3). In addition, in processing of raw products no growth or sporulation of *B. subtilis* is expected to occur.